

Colorado Dairy News

Cooperative Extension, Colorado State University, Fort Collins, Colorado 80523 Volume 10, Number 1
a publication from Integrated Livestock Management (www.combs.colostate.edu/ilm/)

Embryonic Loss:

*What causes it, what amount is normal,
and how do I manage it?*

Kevin McSweeney, DVM, ILM, Dairy Specialist

How frustrating is it to have a cow diagnosed pregnant at the 45-day vet check and then discover at dry check (day 220) that she has lost the fetus somewhere along the way? The precise dollar loss, of course, will vary with milk prices, location, and management approaches, but as an example some research has estimated the cost of every day open in four time intervals:

- Open between day 100 and 130: \$0.50 per day
- Open between day 130 to 160: \$1.42 per day
- Open between day 160 to 175: \$2.99 per day
- Open after day 175: \$4.52 per day

Fertilization rates do not appear to have changed significantly in the last 20 years, and increased embryonic loss is assumed to be a significant cause of poorer calving rates and the overall decline in dairy cow fertility. In addition to the apparent increase in rate of embryonic loss, cows that have lost their fetus are not showing estrous as clearly or estrous is not being recognized as readily resulting in "open" cows at the dry check; an often overlooked cause of economic loss.

Pregnancy diagnosis by ultrasound (~day 26-30) can be performed two weeks earlier than diagnosis by rectal palpation. Early embryonic death (EED) can be diagnosed accurately and open cows rebred immediately resulting in cost savings.

What is the normal rate of embryonic loss? Some level of embryonic loss is normal because it represents the loss of unhealthy embryos. To estimate the normal rate of embryonic death, studies of timed breeding protocols can be reviewed. These studies using flushed embryos estimate fertilization rates among lactating dairy cattle to be between 66-87%. This wide span is due to the timing of insemination used in different study protocols: breeding closer to the last GnRH dose will result in a lower fertilization rate but increase the percentage of high quality embryos. When reviewing these types of studies as a whole, the majority of early embryonic death occurs prior to day 27. Affected cows will most often be seen as rebreeds. Current information obtained by using reproductive ultrasound on large expanding dairies have found 10-20% embryonic loss from day 27 to day 45.

What are some normal EED rates in Colorado dairies? I have been working closely with two dairies here in eastern Colorado. Table one illustrates early pregnancy loss data collected with the use of reproductive ultrasound and rectal palpation.

(Please continue on page 3, under EED)

Colorado State University and U.S. Department of Agriculture cooperating. Cooperative Extension programs are available to all without discrimination.

**BSE in the U.S:
What do you do
if a reporter calls
you?**

Identification of a dairy cow in the U.S. with BSE is old news, but the livestock industry will be dealing with the inevitable publicity for quite awhile. Included in this issue as an insert is some helpful information from the National Cattlemen's Beef Association (NCBA) on BSE. Also, please call Bill Wailes, CSU Extension Dairy Specialist, and Frank Garry, Integrated Livestock Management Coordinator, if you have further questions.

Because you own cattle, there is a good possibility that you may be contacted by the media to comment on any aspect that evolves out of the continuing BSE story. It can be very difficult to make your position known clearly. How should producers work with reporters to ensure that accurate and positive information reaches consumers? Below are five basic guidelines for talking to the media that apply at anytime but are particularly helpful when discussing a "hot" topic like BSE. This information is provided by the NCBA.

1. Screen media calls: Ask who the reporter is, who they work for, what they want to talk about, when the story *(Please continue on pg 4, under BSE)*

Table of Contents

ILM Student Seminar	3
BSE in U.S.	Insert
Milking Procedures	Insert

Important Dates:
Mark Your Calendar

January 27, 2004: Colorado Dairy Nutrition Conference; Greeley, Colorado Contact: William Wailes; Dairy Specialist (970)491-5390 w.wailes@colostate.edu

January 28 & 29, 2004: Colorado Dairy Days;Greeley, Colorado: Contact Keith Maxey, Extension Agent Weld County (970)356-4000 ext 4475 Arlan Marrs (303) 659-8682.

February 9, 2004: Colorado Water Quality Control Commission Rulemaking Hearing for CAFO's. Colorado Department of Health, Denver Colorado; For more information contact: Meg Collins (720)201-2870 or megcollins2@aol.com



**Integrated
Livestock
Management**

Colorado Dairy News
is published

bimonthly as a service to those people interested in the health and welfare of the Colorado dairy industry.

Past issues are available on the ILM website

www.combs.colostate.edu/ilm

For further information, contact Dr. Ragan Adams, Editor, ILM, CSU-VTH 300 West Drake Rd Ft Collins, CO 80523

(970)297-0371;

radams@lamar.colostate.edu

Material published in the Colorado Dairy News is not subject to copyright. Permission is therefore granted to reproduce articles although acknowledgement of the source is requested.

A Message From Your Extension Dairy Specialist.....

CAFO Hearings: The Colorado Water Quality Division has conducted several stakeholders meeting with representatives of the livestock industry and others to assist the Division in the development of regulatory language, which will bring the state CAFO regulations in line with new federal requirements. These meetings have reviewed the areas where the state has flexibility within the federal regulations. At the same time, to assist livestock producers, the Division developed and distributed a Notice of Intent (NOI), which allows state CAFO's to inform the Division of the adequacy of their storage and other steps that producers are taking to comply with the state's permit requirements. This whole process will come to its final decision-making on rules that will affect you and your dairy operation's forever. If you have any questions or any misunderstandings about this process please talk with our representatives that have been involved with and have been representing you at these meetings.

MILC Payments:The Milk Income Loss Contract payments will begin in January 2004. If you have any question contact your local Farm Service Agency office in your county.

COOL Delay: Lawmakers and the White House officials, negotiating details of an overdue government-spending bill, have agreed to delay country-of-origin labeling for two more years. The labeling law was imposed in the 2002 Farm Bill, with mandatory labeling set to begin in September 2004. The two-year delay on COOL applies to meats, produce and farm-raised fish. Producers in Colorado will hear more on this issue as the industry makes rules for you to operate under for identification.

William R. Wailes, Colorado Extension Dairy Specialist

Commodity Price Quotes

By-Product Feeds	Price/Ton Spot Loads	Price/Ton Jan/Feb/March 2004
Bakery Waste	\$102.00	NQ
Blood Meal	\$775.00	NQ
Corn Gluten Feed	\$130.00	\$125.00
Corn Hominy	\$98.00	\$98.00
Flaked Corn	\$6.00/cwt	NQ
Whole Corn	\$5.20/cwt	NQ
Whole Cottonseed	\$189.00	\$195.00
Distillers Grains	\$148.00	\$148.00
Pork - Meat & Bone Meal	\$315.00	NQ
Tallow	\$0.25 /lb.	NQ
SBM - 48%	\$250.00	\$253.00
Wheat Middlings	\$105.00	\$95.00
Soybean Hulls	\$112.00	\$110.00
Canola Meal	\$185.00	\$185.00

These price quotes are delivery at Greeley, Co

(EED, continued from page 1)

Table One: Embryonic loss on two Colorado dairies

Dairy A milks 500 cows and had a total embryonic loss from day 28 to day 100 of 14%: 8% embryonic loss occurred from day 28 to 35, 5% from day 35 to day 55, 1% loss from day 55 to 100.
Dairy B milks 1200 cows and had a total embryonic loss from day 28 to 95 of 14.3%: 10.1% embryonic loss occurred from day 28 to day 60. 4.2% loss from day 60 to 95.

There are many reasons for embryonic loss. The following factors should be considered if the embryonic loss rate at a dairy is not acceptable.

The oocyte: The oocyte or unfertilized egg develops within the follicle. Studies suggest that once this follicle and oocyte attain dominance (usually when follicles reach ~10mm in diameter) after a few days the quality of this oocyte starts to decline. It is quite common for lactating dairy cows to ovulate poor quality oocytes, and one reason may be follicles that persist contain poor quality oocytes. These inferior oocytes lead to inferior quality embryos which are more apt to be lost.

Uterine environment also contributes to embryonic loss. To evaluate the effect of the maternal uterine environment on EED, in one study day 7 embryos were flushed from non-lactating Holstein heifers and lactating Holstein cows with poor fertility. Only high quality embryos were used. The lactating cows received the embryos from the heifers, and the heifers received the embryos from the cows. The heifers attained pregnancy rates very similar to rates attained using embryos from other heifers, but the cows had pregnancy rates similar to rates attained using their own embryos. By neutralizing the difference in origin of embryos, this study suggests that the poor pregnancy rate in the cows was due to uterine environment rather than poor quality embryos.

Blood progesterone levels must be maintained to support pregnancy. Lactating dairy cattle have been shown to have lower blood progesterone levels than they did 30 years ago. Although the cause of low blood progesterone is not completely understood one well accepted hypothesis is that it results from the industry's efforts to maximize feed intake and boost milk production. When feed intake increases, more blood passes through the liver which breaks down the progesterone and estrogen. This decrease in progesterone levels is thought to lead to more embryonic loss.

If lower blood progesterone levels are detrimental to embryonic health, what can be done to increase progesterone levels? The corpus luteum (CL) located on the ovary is the main source of progesterone. If the numbers of CL's are increased, blood progesterone levels should also increase. One way to accomplish increasing the number of CL's is to give Human Chorionic Gonadatropin (HCG) around day 5 post breeding. This causes the next follicular wave to ovulate and creates additional CL's, thus increasing blood progesterone levels. One study on a large commercial dairy in California during cool weather months showed nearly a 40% increase in conception rates (34.2% CR in the non-treated group versus 47.8% CR in the HCG treated group) by giving HCG day 5 post A.I. 86.2% of HCG treated cows had accessory CL's and their blood progesterone levels were increased by 5.0 ng/mL (a big increase) when compared to controls.

(Please continue on page 4, under EED)

ILM Summer 2003 Student Forum

The ILM Summer Student Program involves students in ILM project activities during the summer academic break. It is designed to give the students experience and insight into livestock health problem solving. Each student is assigned a faculty mentor who oversees a project, generally concerning a livestock producer problem. The students also attend weekly seminars conducted by ILM faculty and participate in field activities outside of their own project area to broaden their experience. The program, begun in the summer of 2000 as a unique opportunity for veterinary students at Colorado State University, is supported by donations from livestock industry companies and private livestock producers.

Ten students involved in activities during the 2003 summer break presented their research findings December 3rd. Sophomore veterinary students **Jace King** and **Michael Suite** worked with Dr Kevin McSweeney to collect information contained in the lead article on EED in this issue of CDN. **Katie Jones**, a junior veterinary student studied mastitis and udder preparation in dairy cattle with Dr Heather Hirst. Dusti Pruna a sophomore veterinary student worked with Dr John Wenz on a study concerning the use of recycled sand bedding. Sarah Tomlison, a senior veterinary student worked with Dr Jason Lombard on a study of dystocia, calf morbidity and mortality that was reported in the July 2003 CDN and at the annual conference of the American Association of Bovine Practitioners. Senior veterinary student Julie Severidt and Gina Henriksen, another sophomore veterinary student, worked on educational tools for dairy workers involving lameness, hoof care and sick cows.

Students, producers or donors interested in participating in this program can contact Dr Frank Garry, Coordinator of ILM (970) 297-0371.

(EED, continued from page 3)

General maternal health: There are also many disease processes that can cause EED. Such common ones are BVD, coliform mastitis, trichomoniasis, neospora, vibrio, and *Leptospira hardjo*. Leptospirosis as a cause of reproductive failure has been gaining considerable attention lately and there is a new vaccine for its prevention.

How do I manage EED?

EED is a very complex problem, and requires diligence to deal with it. Although it may be minimized, it will never be eliminated. Certain amounts of EED are normal, but we need to manage it to minimize the negative impact on the bottom line. Another period of pregnancy loss also occurs after day 45-50 post breeding. In a recent study Vasconcelos estimated this fetal loss after day 56 to be 9.2%. Most dairies diagnose pregnancy around 40-50 days post-breeding and then reconfirm the pregnancy at dry-off (day 220), but there can be considerable loss occurring between these two checks. If a specific disease (BVD, leptospirosis, etc.) is responsible it needs to be identified and managed.

I recommend determining what level of embryonic loss you are experiencing and also when it is occurring. If you find that you have a considerable proportion of cows open at dry check, you might consider adding an additional pregnancy check somewhere near day 100 post-breeding. For those suspicious of the use of either ultrasound for early pregnancy diagnosis or timed breeding protocols, neither have been shown to increase the amount of EED when compared to non-synched heat bred cattle or those diagnosed pregnant by rectal palpation.

If you are fortunate enough to have ultrasound on your farm, you probably have already been managing EED, with rechecks of all your pregnant cows done around day 40-60 post breeding. There has been some work out of the University of Wisconsin that showed the greatest amount of embryonic loss after day 28 occurred within that next week (day 28-35 post breeding). The data from my study also showed that 55% of the total embryonic loss occurring after day 28 occurred between day 28 and day 35 post breeding. If the vast majority of these embryos are lost during this week, why wait longer to reconfirm. On Dairy A (500 cows) we recheck all pregnant cows one week after the day 28 scan. We reconfirm the remaining pregnancies by rectal palpation near day 55, then again day 100 and finally at dry-off. We start resynchronization on all open cows whenever they are diagnosed and when their ovaries contain a CL (the best time to start resynch programs is when a cow has a CL). Some cows will still have a dead embryo and intrauterine debris and need to be "cleaned-up" by giving a shot of lualyse prior to starting the resynchronization protocol.

This approach may be too aggressive for the average dairy, but some variation of this protocol may be beneficial. Open cows cost money. If too many cows are coming up open at dry-off, you are losing considerable money. Not only are you losing the potential increase in milk from a peak lactation cow, but also with prolonged lactation you may lose the cow. These cows are usually far into their lactation and their milk is often too low to keep them around to try and get them pregnant again. They go to the sale barn where a cow you thought had a value of \$1200-\$1800 prior to the pregnancy check now gets you \$300-\$600 as a cull cow. Identifying embryonic loss in cows as soon as possible and getting affected cows rebred promptly can mean the difference in whether you freshen these cows on your dairy in 9+ months or cull them in a few months.

(BSE, continued from page 1)

deadline is, and when you can call them back. You don't have to talk with reporters the moment they call. They generally understand that you will need to call them back because you are busy, but they do have deadlines and will expect you to get back with them quickly. Use the time before the call back to organize your thoughts, get answers you need, and line up talking points. If you don't feel comfortable talking with them, refer them to your state or national organization. If you do an interview, let your state organization know.

2. Focus on your message: Select three main points you want to get across and repeat them throughout the interview. Try to maintain control of the direction of the interview by returning to your main message. Use the basic technique of briefly answering the reporter's question, even if the answer is a simple "yes" or "no", and then bridge to one of your prepared messages.

3. Never speculate: Focus on the facts and don't allow reporters to lead you into speculation, conjecture, or places where you don't want to go. It is perfectly acceptable to tell a reporter "I don't know the answer to that, but I will find out and get back to you." Then do get the reporter's contact information and do get back to him or her as soon as possible.

4. Take advantage of the experts: Direct reporters to places they can find reliable information (your local cattle or beef association, NCBA and the Web site at www.bseinfo.org, or the USDA site at www.usda.gov.)

5. Be positive: You can set the tone of the article by your attitude. Avoid repeating negative language and look for ways to bridge the discussion to positive messages like: "The U.S. has the safest beef in the world."

For further information contact NCBA in Centennial, Co at 303-694-0305 or in Washington, DC at 202-347-0228.